

Proof Script for SS18 Convex Optimization Lecture*

Last updated: April 23, 2018

1 Convex Analysis

Theorem 1.1 (separation of convex sets). *Let C_1, C_2 be nonempty convex subsets in \mathbb{E} such that $C_1 \cap C_2 = \emptyset$ and C_1 is open. Then there exists a hyperplane separating C_1 and C_2 , i.e. $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ such that*

$$\langle v, u^1 \rangle \geq \alpha \geq \langle v, u^2 \rangle, \quad \forall u^1 \in C_1, u^2 \in C_2.$$

Proof. (i) Claim: Let $C \subset \mathbb{E}$ be closed, convex set, and $w \in \mathbb{E} \setminus C$. Then $\exists v \in \mathbb{E}, v \neq 0, \alpha \in \mathbb{R}$ s.t. $\langle v, w \rangle > \alpha \geq \langle v, u \rangle \quad \forall u \in C$.

Consider the projection of w onto C , i.e. set $u^* := \arg \min_{u \in C} \frac{1}{2} \|u - w\|^2$ or, equivalently, let $\langle u - u^*, u^* - w \rangle \geq 0 \quad \forall u \in C$.

Now set $v := w - u^* \neq 0$. Then $\forall u \in C$, we have $\langle v, w \rangle = \langle w - u^*, w \rangle = \|w - u^*\|^2 + \langle w - u^*, u^* \rangle \geq \|w - u^*\|^2 + \langle w - u^*, u \rangle = \|v\|^2 + \langle v, u \rangle$. Set $\alpha := \sup\{\langle v, u \rangle : u \in C\}$. Note $\alpha < \infty$ since $\langle v, u \rangle \leq \langle v, u^* \rangle \quad \forall u \in C$. Thus $\langle v, w \rangle > \alpha \geq \langle v, u \rangle \quad \forall u \in C$, which proves the claim.

(ii) Let C_1 be an open, convex subset of \mathbb{E} , and $C_2 = \{\bar{w}\}$ with $\bar{w} \in \mathbb{E} \setminus C_1$. Since $\mathbb{E} \setminus C_1$ is closed, $\exists w^k \in \mathbb{E} \setminus \text{cl } C_1$ s.t. $w^k \rightarrow \bar{w}$. For each w^k , by (i), $\exists v^k \in \mathbb{E}$ with $\|v^k\| \equiv 1$ s.t. $\langle v^k, w^k \rangle \leq \langle v^k, u^1 \rangle \quad \forall u^1 \in C_1 \subset \text{cl } C_1$. Hence $v^k \rightarrow \bar{v} \in \mathbb{E}$ along a subsequence s.t. $\|\bar{v}\| = 1$ and $\langle \bar{v}, \bar{w} \rangle \leq \langle \bar{v}, u^1 \rangle \quad \forall u^1 \in C_1$.

(iii) Consider C_2 as a general convex subset of \mathbb{E} . Set $C := C_2 - C_1 = \{u^2 - u^1 : u^1 \in C_1, u^2 \in C_2\}$. Note that C is a convex, open set, and $0 \notin C$. By (ii), $\exists \bar{v} \in \mathbb{E}$ with $\|\bar{v}\| = 1$ s.t. $\langle -\bar{v}, u^2 - u^1 \rangle \geq \langle -\bar{v}, 0 \rangle = 0$ or, equivalently, $\langle \bar{v}, u^1 \rangle \geq \langle \bar{v}, u^2 \rangle \quad \forall u^1 \in C_1, u^2 \in C_2$. Set $\alpha := \sup\{\langle \bar{v}, u^2 \rangle : u^2 \in C_2\}$, then we conclude that $\langle \bar{v}, u^1 \rangle \geq \alpha \geq \langle \bar{v}, u^2 \rangle \quad \forall u^1 \in C_1, u^2 \in C_2$. \square

Theorem 1.2. *A proper convex function $J : \mathbb{E} \rightarrow \bar{\mathbb{R}}$ is locally Lipschitz at any $u \in \text{rint dom } J$.*

Proof. Throughout the proof, we consider $J : \text{aff dom } J \rightarrow \bar{\mathbb{R}}$.

(i) Claim: If $M = \sup\{J(v) : v \in B_\epsilon(u)\} < \infty$ with $\epsilon > 0$, then J is locally Lipschitz at u .

First, by convexity of J we have $\forall v \in B_\epsilon(u) : J(v) \geq 2J(u) - J(2u - v) \geq 2J(u) - M$. Thus, $\sup\{|J(v)| : v \in B_\epsilon(u)\} \leq M + 2|J(u)|$.

Next, we show J is Lipschitz on $B_{\epsilon/2}(u)$. Let $v, w \in B_{\epsilon/2}(u)$ be given. Take $z \in B_\epsilon(u)$ s.t. $w = (1 - t)v + tz$ for some $t \in [0, 1]$ and $\|z - v\| \geq \epsilon/2$. By convexity, $J(w) - J(v) \leq t(J(z) - J(v)) \leq 2t(M - J(u))$. Since $t(z - v) = w - v$, we have $t = \|w - v\| / \|z - v\| \leq 2\|w - v\| / \epsilon$ and $J(w) - J(v) \leq (4(M - J(u)) / \epsilon) \|w - v\|$. Analogously, one can show $J(v) - J(w) \leq (4(M - J(u)) / \epsilon) \|w - v\|$. Hence, J is Lipschitz on $B_{\epsilon/2}(u)$ with modulus $4(M - J(u)) / \epsilon$.

*Please report typos to: tao.wu@tum.de

(ii) Let $u \in \text{rint dom } J$ and $n = \dim(\text{aff dom } J)$. Then by Carathéodory's theorem, $\exists \{\alpha^i\}_{i=1}^{n+1} \subset (0, 1)$, $\{u^i\}_{i=1}^{n+1} \subset \text{dom } J$ s.t. $u = \sum_{i=1}^{n+1} \alpha^i u^i$, $\sum_{i=1}^{n+1} \alpha^i = 1$, i.e. u belongs to the interior of the convex hull of $\{u^i\}_{i=1}^{n+1}$. Thus one can apply (i) to assert that J is locally Lipschitz at u . \square

Theorem 1.3. *For any proper convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, if $u^* \in \text{dom } J$ is a local minimizer of J , then it is also a global minimizer.*

Proof. By the definition of a local minimizer, $\exists \epsilon > 0$ s.t. $J(u^*) \leq J(u) \forall u \in B_\epsilon(u^*)$. For the sake of contradiction, assume $\exists \bar{u} \in \mathbb{E}$ s.t. $J(\bar{u}) < J(u^*)$. By convexity of J , we have $J(\alpha \bar{u} + (1 - \alpha)u^*) \leq J(u^*) - \alpha(J(u^*) - J(\bar{u})) < J(u^*) \forall \alpha \in (0, 1]$. This violates the local optimality of u^* as $\alpha \rightarrow 0^+$. \square

Theorem 1.4. *Any proper function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$, which is bounded from below, coercive, and lsc, has a (global) minimizer.*

Proof. Let $\{u^k\}$ be an infimizing sequence for J , i.e. $\lim_{k \rightarrow \infty} J(u^k) = \inf_{u \in \mathbb{E}} J(u) > -\infty$. Since $\{J(u^k)\}$ is uniformly bounded from above, by coercivity of J , $\{u^k\}$ is uniformly bounded. By compactness, $u^k \rightarrow u^*$ along a subsequence. Since J is lsc, we have $J(u^*) \leq \liminf_{k \rightarrow \infty} J(u^k) = \inf_{u \in \mathbb{E}} J(u)$, which implies $J(u^*) = \inf_{u \in \mathbb{E}} J(u)$ or u^* is a minimizer of J . \square

Theorem 1.5. *The minimizer of a strictly convex function $J : \mathbb{E} \rightarrow \overline{\mathbb{R}}$ is unique.*

Proof. Let $u, v \in \mathbb{E}$ be two (global) minimizers s.t. $u \neq v$ and $J(u) = J(v) = J^*$. By strict convexity of J , $J(\alpha u + (1 - \alpha)v) < \alpha J(u) + (1 - \alpha)J(v) = J^*$ for all $\alpha \in (0, 1)$, which contradicts the global optimality of u and v . \square